



Pre-made Target Over-expression Lentivirus

CAT#	Product	Accession ID
LVPxxx	Concentrated Expression Lentivirus (200ul at titer of 1×10^8 IFU/ml)	

Amount: 200ul/vial (1×10^8 IFU/ml)/each.

Storage: -70°C , avoid repeat freeze/thaw cycles. Stable for 6 months at -70°C .

Product Description:

GenTarget's Lentivector system is Human Immunodeficiency Virus-1 (HIV) based lentivector plasmids for gene expression and knockdown. The lentivectors are used to generate lentiviral particles (lentivirus) that can be transduced into most mammalian cell types, including stem cells, primary cells, and non-dividing cells both *in vivo* and *in vitro*. Lentiviral Particles stably integrate into the transduced cells' genome for long term expression, making lentivirus a great gene transfer agent.

This target over-expression Lentivirus products, is generated from GenTarget's Lentivector. The target is expressed under the enhance EF1a promoter which is less cell type dependent and not be silenced *In vitro* and *In vivo*, also strong in many primary cells and T cells, B cells, and more. The lentivirus contains the **Puromycin** selection under RSV promoter.

The Lentivector is adapted the most advanced biosafety features, including the self-inactivation feature in its 3' LTR, and only generates the replication-incompetent lentivirus. See the lentivector's map scheme below.



The [Negative Control Lentivirus](#) can be used to establish the controls for lentivirus treatment in a given cell line. The control lentivirus, CAT# [EF1a-Null-Puro](#), has the identical lentivector backbone with the puromycin resistance, as that of target expression, but does not express any target. Please see also "[FAQs about premade lentiviral particles](#)".



Key features:

- 1) High target expression levels driven by strong enhanced EF1a promoter, best for primary cells, suspension cells, T cells, B cells and more;
- 2) Puromycin killing selection;
- 3) Ready to use and easy to use: simply add it into your cell culture, No need any other reagents or procedures;

Transduction Protocols:

1) Transduction Protocol for Adhesive cells :

Note: lentivirus is provided as ready to use. Simply add it into your cell culture; the amount to add depends on cell type.

Quick transduction protocol: add 50 μ l of virus into each well of 24-well-plate where cell density is 50% to 75%. After 72 hours (no need to change medium), check transduction rate by fluorescence signal. For stable cell line generation, pass cells into medium containing antibiotic or carry out the fluorescence cell sorting.

Day 0:

Seed cells in complete medium at the appropriate density and incubate overnight.

Note: at the time of transduction, cells should be 50%-75% confluent. For example, seed HeLa cells at $0.5 \times 10^5/\text{ml} \times 0.5\text{ml}$ in a well of a 24-well plate.

Day 1:

- Remove the culture medium and add 0.5ml fresh, warm, complete medium.
- Thaw the lentiviral stock at room temperature and add the appropriate amount of virus stock to obtain the desired MOI, normally use MOI from 2 to 10.
- Return cells to 37°C, CO₂ incubator.

Note: If you do not use all of the virus at one time, you can re-freeze the virus at -80 °C for future use; virus titer will decrease by ~10% for each freeze/thaw cycle.

Day 3:

At ~72hr after transduction, check the transduction rate under fluorescence microscopy or calculate the exact transduction rate by



flow cytometry (FACS or Guava). Carry out the expression detection assay.

Day 3 +

Sort transduced cells by FACS, or select the transduced cell by antibiotic resistance. A pilot experiment should be done to determine the antibiotic's kill curve for your specific cell line (refer to the pertinent literature on generation of stable cell lines).

2) Transduction Protocol for Suspension Cells:

Grow cells in complete suspension culture medium; use a shaking flask in a CO₂ incubator if necessary.

Measure cell density. When density has reached $\sim 3 \times 10^6$ cells/ml, measured viability should be > 90%. Dilute cells into 1×10^6 cell/ml in complete medium.

Day 1:

- Thaw lentiviral particles at room temperature.
- Add premade lentiviral particles into the diluted cells at a ratio of: 50 to 100 μ l virus per 0.5 ml of cells (Note: depending on cell type, you may need to use more or less virus).
- Grow cells in a shaking flask in a CO₂ incubator.

Day 2:

At 24 hours after transduction, add an equal amount of fresh medium containing relevant antibiotics. **Note:** amount of antibiotic depends on cell type. Continue growing cells in CO₂ incubator.

Day 3:

At 72 hours after transduction, check fluorescence with a fluorescence microscope or calculate the transduction efficiency using a cell sorter such as FACS or Guava. Sort for fluorescence positive cells and maintain antibiotic selection to generate a stable cell line.

Safety Precaution:

Gentarget lentiviral particles adapt must advanced lentiviral safety features (using the third generation vectors with self-inactivation SIN-3UTR), and the premade lentivirus is replication incompetent. However, please use extra caution when using lentiviral particles. Use the lentiviral particles in Bio-safety II cabinet. Wear glove all the time when handling Lentiviral particles!



Please refer CDC and NIH's guidelines for more details regarding to safety issues.

References:

1. J Virol. 2000 November; 74(22): 10778–10784.
2. Hum Gene Ther (2003) 14: 1089-105.
3. Mol Ther (2002) 6: 162-8.
4. NIH Guidelines for [Biosafety Considerations for Research with Lentiviral Vectors](#). (Link).

Warranty:

This product is for research use only. It is warranted to meet its quality as described when used in accordance with its instructions. GenTarget disclaims any implied warranty of this product for particular application. In no event shall GenTarget be liable for any incidental or consequential damages in connection with the products. GenTarget's sole remedy for breach of this warranty should be, at GenTarget's option, to replace the products.

Attachment: GenTarget's pre-made lentivirus product categories.

Product Category	Product Description (please click into each category's page)
Pathway Reporter	Repoter Lentivirus for all kinds of pathway screening assays
Cell Immortalization	Lentivirus for cell immortalization: Large T-antigen, hTERT, EBNA1/EBNA2, HpV16-E6/E7, Adenovial E1A, Kras_G12V, HOXA9, et al.
ImmunoOncology Research	Lentivirus products for immuno therapy research: CAR and TCR; Assay Cell Lines for T-cell targeted killing assay and other cell-based assays; over-expression lentivirus products for the immune response targets; Cell surface antigens (CDs); immune checkpoint / Receptors; CRISPR gene Repair and knock-IN lentivirus; CRISPR knockout lentivirus;
CAR-T, TCR Lentivirus	CARs Lentivirus: Anti-CD19 /CD20 /CD22 /BCMA /hHER2 /HLA-A2 /TGFβ; TCRs : MART-1/ NY-ESO1/ CD1d-α-GalCer/ TRαV3-F2A-TRβV5-6;
CRISPR Gene Editing	Preamde lentivirus express humanized wild-type Cas9 endonuclease, the dCas9 , gRNAs, CRISPR gene editing research



Product Category	Product Description (please click into each category's page)
Epigenomic: CRISPRi and CRISPRa	" dCas9-Protein " fusion Lentivirus for epigenomic modification, resulted in CRISPR interference (CRISPRi) or activation (CRISPRa).
Cell-Specific Reporter	a set of reporter lentiviruses to express a luminescence or fluorescent reporter (firefly Luciferase, Renilla luciferase, RFP or GFP fluorescent marker) under a tissue specific promoter
Infectious Antigens	Lentivirus that express all kinds of infectious antigens with C-term 6His-tag.
Virus Like Particles (VLP)	Lentiviral Like Particles, pseudo-typed with a different envelope proteins.
Non-integrating LV	Integration Defective Lentivirus, express different targets for transient expression without the unwanted insertional mutagenesis.
shRNA Knockdown	Knockdown verified and customized shRNA lentivirus for target knockdown,
microRNA lentivirus	Premade lentivirus expression human or mouse precursor miRNA . And anti-miRNA lentivector and virus for human and mouse miRNA.
Anti-miRNA lentivirus	Pre-made lentivirus expression a specific anti-miRNA cassette.
Human and mouse ORFs	Premade lentivirus expressing a human, mouse or rat gene with RFP-Blasticidin fusion dual markers.
Luciferase expression	Premade lentivirus for all kinds of luciferase protein expression: firefly and Renilla, Red-Luc and more , with different antibiotic selection markers.
Fluorescent Markers	Lentivirus express all commonly used fluorescent proteins: GFP, RFP, CFP, BFP YFP, mRFP, unstable GFP and others.
Luminescent Imaging	Lentivirus express Nano-Lantern as Bio-probes for in vivo imaging of sub-cellular structural organization and dynamic processes in living cells and organisms
Sub-cellular Imaging	Lentivirus contain a well-defined organelle targeting signal fused to a fluorescent protein, great tools for live-cell imaging and for dynamic investigation of sub-cellular signal pathways.



Product Category	Product Description (please click into each category's page)
Cytoskeleton Imaging	A fluorescent marker (GFP, RFP or CFP) fusion with a cellular structure protein, provides a convenient tool for visualization of cytoskeletal structure
Unstable GFP	Lentivirus express the the destabilized GFP (uGFP) which provides fast turnover responses in signal pathway assay and in knockdown / knockout detection
near-infrared RFP	The near-infrared Red fluorescent (niRFP) expression Lentiviruses provides the whole-body images with better contrast and brighter images
Fluorescent-ORF fusion	Pre-made lentivirus expression a " GFP/RFP/CFP-ORF " fusion target.
CRE recombinase	Premade lentivirus for expressing nuclear permeant CRE recombinase with different fluorescent and antibiotic markers.
CRE, Flp ColorSwitch	Lentivirus expressing "LoxP-GFP-Stop-LoxP-RFP" or "FRT-GFP-Stop-FRT-RFP" cassette, used to monitor the CRE or Flp recombination event in vivo.
SEAP Reporter	lentivirus expressing SEAP under different promoters (TetCMV, EF1a, CAG, Ubc, mPGK, Actin-beta or a signal pathway responsive promoter),
TetR Repressor	Premade lentivirus expressin TetR (tetracycline regulator) protein, the repressor protein for the inducible expression system.
rtTA Expression	rtTA binds to the tetracycline operator element (TetO) in the presence of doxycycline (Dox). Used for Tet-On /OFF inducible system.
iPS factors	Premade lentivirus for human and mouse iPS (Myc, NANOG, OCT4, SOX2, FGF4) factors with different fluorescent and antibiotic markers
LacZ expression	Express different full length β-galactosidase (lacZ) with different selection markers
Negative control lentiviruses	Premade negative control lentivirus with different markers : serves as the negative control of lentivirus treatment, for validation of the specificity of any lentivirus target expression effects.
Other Enzyme expression	Ready-to-use lentivirus, expressing a specific enzymes with different selection markers.
Ultra titer	Ultra-titer lentivirus used for the hard-to-transduced



GenTarget Inc

7930 Arjons Drive, Suite B
San Diego, CA 92126, USA
Phone: 1 (858) 265-6446
Fax: 1 (800) 380-4198
Email: orders@gentarget.com

Product Category	Product Description (please click into each category's page)
lentivirus	cells and for in vivo manipulation of sperm cells, or stem cells.