



Human hTERT Expression Lentivirus for Cell Immortalization

CAT#	Product Name	Amounts
LVP1130-Neo	hTERT (CMV, Neo) lentivirus	200ul, (1 x 10 ⁷ IFU/mL) in DMEM medium with premixed Polybrene
LVP1130-Bsd	hTERT (CMV, Bsd) lentivirus	
LVP1130-Puro	hTERT (CMV, Puro) lentivirus	
LVP1130-Zeo	hTERT (CMV, Zeo) Lentivirus	
LVP1130-Hygro	hTERT (CMV, Hygro) Lentivirus	
LVP1130-GB	hTERT (CMV, GFP-Bsd) lentivirus	
LVP1130-GP	hTERT (CMV, GFP-Puro) lentivirus	
LVP1130-RB	hTERT (CMV, RFP-Bsd) lentivirus	
LVP1130-RP	hTERT (CMV, RFP-Puro) lentivirus	
LVP1131-Neo	hTERT (EF1a, Neo) lentivirus	
LVP1131-Bsd	hTERT (EF1a, Bsd) lentivirus	
LVP1131-Puro	hTERT (EF1a, Puro) lentivirus	
LVP1131-Zeo	hTERT (EF1a, Zeo) Lentivirus	
LVP1131-Hygro	hTERT (EF1a, Hygro) Lentivirus	
LVP1131-GB	hTERT (EF1a, GFP-Bsd) lentivirus	
LVP1131-GP	hTERT (EF1a, GFP-Puro) lentivirus	
LVP1131-RB	hTERT (EF1a, RFP-Bsd) lentivirus	
LVP1131-RP	hTERT (EF1a, RFP-Puro) lentivirus	200ul, (1 x 10 ⁸ IFU/mL) in PBS solution with premixed Polybrene
LVP1130-Neo-PBS	hTERT (CMV, Neo) lentivirus in PBS	
LVP1130-Bsd-PBS	hTERT (CMV, Bsd) lentivirus in PBS	
LVP1130-Puro-PBS	hTERT (CMV, Puro) lentivirus in PBS	
LVP1130-Zeo-PBS	hTERT (CMV, Zeo) lentivirus in PBS	
LVP1130-Hygro-PBS	hTERT (CMV, Hygro) Lentivirus in PBS	
LVP1130-GB-PBS	hTERT (CMV, GFP-Bsd) lentivirus in PBS	
LVP1130-GP-PBS	hTERT (CMV, GFP-Puro) lentivirus in PBS	



LVP1130-RB-PBS	hTERT (CMV, RFP-Bsd) lentivirus in PBS	
LVP1130-RP-PBS	hTERT (CMV, RFP-Puro) lentivirus in PBS	
LVP1131-Neo-PBS	hTERT (EF1a, Neo) lentivirus in PBS	
LVP1131-Bsd-PBS	hTERT (EF1a, Bsd) lentivirus in PBS	
LVP1131-Puro-PBS	hTERT (EF1a, Puro) lentivirus in PBS	
LVP1131-Zeo-PBS	hTERT (EF1a, Zeo) Lentivirus in PBS	
LVP1131-Hygro-PBS	hTERT (EF1a, Hygro) Lentivirus in PBS	
LVP1131-GB-PBS	hTERT (EF1a, GFP-Bsd) lentivirus in PBS	
LVP1131-GP-PBS	hTERT (EF1a, GFP-Puro) lentivirus in PBS	
LVP1131-RB-PBS	hTERT (EF1a, RFP-Bsd) lentivirus in PBS	
LVP1131-RP-PBS	hTERT (EF1a, RFP-Puro) lentivirus in PBS	
LVP1131	hTERT (EF1a) lentivirus (note: no any antibiotic selection)	200ul (1x10 ⁷ IFU/ml)
LVP1131-PBS	hTERT (EF1a) lentivirus in PBS (note: no any antibiotic selection)	200ul (1x10 ⁸ IFU/ml)

Storage: <-70 °C, avoid repeat freeze/thaw cycles, stable for > 6 months.

Product Description:


Lentiviral particles or lentivirus is a gene delivery tool produced from lentivectors for gene expression or knockdown. GenTarget’s lentivector system is Human Immunodeficiency Virus-1 (HIV) based plasmids for gene expression and knockdown. The lentivectors are used to generate lentiviral particles (lentivirus) that can be transduced into almost all kinds of mammalian cells, including stem cells, primary cells, and non-dividing cells both *in vivo* and *in vitro*. Lentivirus can stably integrate into the transduced cells’ genome for long term expression, making it a great gene transfer agent.



Human Telomerase Reverse Transcriptase (hTERT) plays a role in cellular senescence and also participates in chromosomal repair. When hTERT is exogenously expressed, the cells are able to maintain telomere lengths to avoid cell senescence. Therefore, hTERT is used for primary cell immortalization for variety of cell types. For some primary cell types, the cell immortalization may requires a combination of immortalization method, like, the over-expression of both hTERT and SV40 Large T antigen. (Note: for some cell types, the over-expression hTERT may be toxic, causing cell death. If so, you have to other method, like use SV40 T antigen for the immortalization.)

hTERT expression lentivirus products are generated from GenTarget's re-engineered lentivector system. The hTERT longest transcript (variant 1) of hTERT codon sequence ([NM_198253](#)), was expressed under an enhanced **CMV** (suCMV) or enhanced **EF1a** promoter. The **suCMV promoter** demonstrate the strongest expression in most cell types and the **enhanced EF1a promoter** is active in almost all cell types and less likely to be silenced during long-term culture.

Each Lentivirus is featured with a selection marker (**Neomycin, Puromycin, Blasticidin, Zeocin, Hygromycin**), or an antibiotic-fluorescent fusion dual maker (**RFP-Bsd, RFP-Puro, GFP-Bsd, GFP-Puro**). (see **vector map scheme** below). (**Note**: we also made a product, CAT#: **LVP1131**, that does not contain any selection marker).



VSV-G pseudo-typed lentivirus are generated in 293T cell, and provided as 200 ul aliquots in two formats:



- 1) in DMEM medium containing 10% and 10x Polybrene (60 ug/ml) at titer of 1×10^7 IFU/ml;
- 2) in PBS solution at titer of 1×10^8 IFU/ml, for usage in serum-free cell culture;

For general questions about our ready-to-use particles, please see [FAQ for pre-made lentiviral particles](#) (.pdf) on our website.
(<http://www.gentarget.com/pdf/FAQ-Premade-Lentiviral-particles.pdf>).

Key features:

1. Each lentiviral particles contains a specific antibiotic resistant marker, or **antibiotic-fluorescent** fusion dual marker, used for selecting the transduced cells or generating stable cell lines by antibiotics selection or via fluorescent cell sorting. (or No any selection, CAT#: **LVP1131**).
2. The strongest [suCMV promoter](#) for high expression.
3. The enhance **EF1a promoter** is active in all cell types and do not be silenced during long-term culture.
4. The lentivirus is ready and easy to use, simply add 50ul into one well culture in 24-well plate. No need any other reagents at application.

Transduction Protocols:

1. Transduction Protocol for Adhesive cells:

Note: Pre-made lentivirus is provided ready to use, so it can be simply added into your cell culture; the amount of virus to add depends on cell type. For quick transduction, add 50 μ l of virus into each well of 24-well-plate where cell density is 50% to 75%. After 72 hours (no need to change medium), visualize positive transduction rate by fluorescence microscopy. For stable cell line generation, pass cells into medium containing antibiotic or perform fluorescence cell sorting followed by antibiotic selection.

Day 0:

Seed cells in complete medium at the appropriate density and incubate overnight.

Note: at the time of transduction, cells should be 50%-75% confluent. For example, seed HeLa cells at 0.5×10^5 /ml x 0.5ml in a well of a 24-well plate.



Day 1:

- Thaw the pre-made lentiviral stock at room temperature and add the appropriate amount of virus stock to obtain the desired MOI.
- Return cells to 37°C, CO₂ incubator. Do nothing.
Note: Try to avoid freezing and thawing. If you do not use all of the virus at one time, you may re-freeze the virus at -80 °C for future use; virus titer will decrease by ~10% for each freeze/thaw cycle.

Day 3:

At 48hr~72hr (Depend upon cell type) after transduction, check the transduction rate by fluorescence microscopy or calculate the exact transduction rate by flow cytometry (FACS or Guava).

Day 3 + (optional):

Sort transduced cells by FACS, and select for antibiotic resistance. A pilot experiment should be done to determine the antibiotic's kill curve for your specific cell line (refer to the pertinent literature on generation of stable cell lines).

2. Transduction Protocol for Suspension Cells:

Grow cells in complete suspension culture medium; use a shaking flask in a CO₂ incubator if necessary.

Measure cell density (not grow over 3 million /ml), measured viability should be > 90%. Dilute cells into 1 x 10⁶ cell/ml in complete medium.

Day 1:

- Thaw lentiviral particles at room temperature.
- Add premade lentiviral particles into the diluted cells at a ratio of: 50 to 100 µl virus per 0.5 ml of cells (Note: depending on cell type, you may need to use more or less virus).
- Grow cells in a shaking flask in a CO₂ incubator.

Day 2:

At 24 hours after transduction, add an equal amount of fresh medium containing. Continue growing cells in CO₂ incubator.

Day 3+:

At 48 hour to 72 hours (Depend upon cell type) after transduction, check fluorescence with a fluorescence microscope or calculate the transduction efficiency using a cell sorter such as FACS or Guava. Pass



cells into 0.5 million/ml density in completed medium containing the corresponding antibiotic (**Note:** amount of antibiotic depends on cell type. A killing curve must pre-established). Sort for fluorescence positive cells and maintain antibiotic selection to generate a stable cell line.

Note: Filter wavelength settings:

BFP filter: ~Ex380 ~Em460;
GFP filter: ~Ex450-490 ~Em525;
RFP filter: ~Ex558 ~Em583;

Safety Precaution:

GenTarget lentiviral particles adapts must advanced lentiviral safety features (using the third-generation vectors with self-inactivation SIN-3UTR), and the premade lentivirus is replication incompetent. However, please use extra caution when using lentiviral particles. Use the lentiviral particles in Bio-safety II cabinet. Wear glove all the time when handling Lentiviral particles! Please refer CDC and NIH's guidelines for more details regarding to safety issues.

References:

1. Human Molecular Genetics, Volume 8, Issue 1, 1 January 1999, Pages 137-142.
2. Current Opinion in Genetics & Development. Volume 9, Issue 1, 1 February 1999, Pages 97-103.
3. Carcinogenesis, Volume 26, Issue 5, 1 May 2005, Pages 867-874

Warranty:

This product is for research use only. It is warranted to meet its quality as described when used in accordance with its instructions. GenTarget disclaims any implied warranty of this product for application. In no event shall GenTarget be liable for any incidental or consequential damages in connection with the products. GenTarget's sole remedy for breach of this warranty should be, at GenTarget's option, to replace the products.

Attachment: GenTarget's pre-made lentivirus product categories.

Product Category	Product Description (please click into each category's page)
Pathway Reporter	Lentivirus for all kinds of pathway assays
Cell Immortalization	Lentivirus for cell immortalization: Large T-antigen, hTERT, EBNA1/EBNA2, HpV16-E6/E7, Adenovial E1A, Kras_G12V, HOXA9, et al.



ImmunoOncology Research	Lentivirus products for immuno therapy research, CAR-T, TCR-T, Assay cell lines, and Cell Antigens & Receptors.
CRISPR Gene Editing	Preamde lentivirus express humanized wild-type Cas9 endonuclease, the dCas9 , gRNAs, CRISPR gene editing research
Cell-Specific Reporter	a set of reporter lentiviruses to express a luminescence or fluorescent reporter (firefly Luciferase, Renilla luciferase, RFP or GFP fluorescent marker) under a tissue specific promoter
Infectious Antigens	Lentivirus that express all kinds of infectious antigens with C-term 6His-tag.
Virus Like Particles (VLP)	Lentiviral Like Particles, pseudo-typed with a different envelope proteins.
Non-integrating LV	Integration Defective Lentivirus, express different targets for transient expression without the unwanted insertional mutagenesis.
shRNA Knockdown	Knockdown verified and customized shRNA lentivirus for target knockdown,
microRNA lentivirus	Premade lentivirus expression human or mouse precursor miRNA . And anti-miRNA lentivector and virus for human and mouse miRNA.
Anti-miNA lentivirus	Pre-made lentivirus expression a specific anti-miRNA cassette.
Human and mouse ORFs	Premade lentivirus expressing a human, mouse or rat gene with RFP-Blasticidin fusion dual markers.
Luciferase expression	Premade lentivirus for all kinds of luciferase protein expression: firefly and Renilla, Red-Luc and more , with different antibiotic selection markers.
Fluorescent Markers	Lentivirus express all commonly used fluorescent proteins: GFP, RFP, CFP, BFP YFP, mRFP, unstable GFP and others.
Luminescent Imaging	Lentivirus express Nano-Latern as Bio-probes for in vivo imaging of sub-cellular structural organization and dynamic processes in living cells and organisms
Cytoskeleton Imaging	A fluorescent marker (GFP, RFP or CFP) fusion with a cellular structure protein, provides a convenient tool for visualization of cytoskeletal structure



Unstable GFP	Lentivirus express the the destabilized GFP (uGFP) which provides fast turnover responses in signal pathway assay and in knockdown / knockout detection
near-infrared RFP	The near-infrared Red fluorescent (niRFP) expression Lentiviurs provides the whole-body images with better contrast and brighter images
Fluorescent-ORF fusion	Pre-made lentivirus expression a " GFP/RFP/CFP-ORF " fusion target.
CRE recombinase	Premade lentivirus for expressing nuclear permeant CRE recombinase with different flurescent and antibiotic markers.
LoxP ColorSwitch	Premade lentivirus expressing "LoxP-GFP-Stop-LoxP-RFP" cassette, used to monitor the CRE recombination event in vivo.
SEAP Reporter	lentivirus expressing SEAP under different promoters (TetCMV, EF1a, CAG, Ubc, mPGK, Actin-beta or a signal pathway responsive promoter),
TetR Repressor	Premade lentivirus expressin TetR (tetracycline regulator) protein, the repressor protein for the inducible expression system.
rtTA Expression	rtTA binds to the tetracycline operator element (TetO) in the presence of doxycycline (Dox). Used for Tet-On /OFF inducible system.
iPS factors	Premde lentivirus for human and mouse iPS (Myc, NANOG, OCT4, SOX2, FLF4) factors with different fluorescent and antibiotic markers
LacZ expression	Express different full length β-galactosidase (lacZ) with different selection markers
Negative control lentiviruses	Premade negative control lentivirus with different markers : serves as the negative control of lentiviruses treatment, for validation of the specificity of any lentivirus target expression effects.
Other Enzyme expression	Ready-to-use lentivirus, expressing a specific enzymes with different selection markers.
Ultra titer lentivirus	Ultra-titer lentivirus used for the hard-to-transduced cells and for in vivo manipulation of sperm cells, or stem cells.